



# Emergence of Earliest Continental Landmass

## Why in News

A new study has suggested that the **earliest continental landmass emerged 3.2 billion year ago instead of 2.5 billion years ago** (as per the continental drift theory).


- The study was conducted by researchers from **India, Australia and the US**.

## Continental Drift Theory

- Continental drift theory **deals with the distribution of the oceans and the continents**. It was first suggested by a German meteorologist, **Alfred Wegener in 1912**.
- According to the theory, **all the continents formed a single continental mass- Pangea** and mega ocean- Panthalassa surrounded it.
- Around **200 million years ago Pangea started splitting** and broke down into two large continental masses as **Laurasia and Gondwanaland** forming the northern and southern components respectively.
- Subsequently, **Laurasia and Gondwanaland continued to break into various smaller continents** that exist today.

//





## Key Points

### ▪ About:

- The study has **challenged** the widely accepted view that the **continents rose from the oceans about 2.5 billion years ago.**
- It suggests **this happened 700 million years earlier — about 3.2 billion years ago —** and that the **earliest continental landmass to emerge may have been Jharkhand's Singhbhum region.**
  - Patches of the earliest continental land, however, exist in Australia and South Africa, too.
  - Geological similarities have linked the Singhbhum craton to cratons in South Africa and Western Australia.

### ▪ Major Findings:

- **River Channels, Tidal Plains and Beaches:**
  - The answer to when the first land masses were formed lay in the sedimentary rocks of the region. **Scientists have found a particular type of sedimentary rocks, called sandstones.**
  - Later on they **found the age by analysing the uranium and lead contents of**

### **tiny minerals.**

- These **rocks were 3.1 billion years old**, and were formed in ancient rivers, beaches, and shallow seas.
- All these water bodies could have only existed if there was continental land. Thus the inference was drawn that the **Singhbhum region was above the ocean before 3.1 billion years ago.**

### ◦ **Extensive Volcanism:**


- The researchers also **studied the granites** that form the continental crust of Singhbhum region.
- These granites are **3.5 to 3.1 billion years old** and formed **through extensive volcanism** that happened about 35-45 km deep inside the Earth and continued on-and-off for hundreds of millions of years until all the magma solidified to form a thick continental crust in the area.
- Due to the thickness and less density, the **continental crust emerged above the surrounding oceanic crust owing to buoyancy** (the quality of being able to float).

### ▪ **Evolution of Organisms:**

- The earliest emergence of continents would have **contributed to a proliferation of photosynthetic organisms**, which would have increased oxygen levels in the atmosphere.
- Weathering of the cratons would have led to nutrient runoff, supplying the ocean with phosphorus and **other building blocks for early life.**
  - **Craton** are the stable interior portion of a continent characteristically composed of ancient crystalline basement rock.

### ▪ **Significance:**

- At a time when the **entire world was debating about changes in climate**, it is very **important to understand how our atmosphere, oceans and climate came into existence** and how they interacted with geological processes operating deep inside Earth to make our planet habitable.
- It will allow us to link the interior of Earth to its exterior in deep time.
  - India has three other ancient continental fragments — **Dharwar, Bastar and Bundelkhand regions.** To understand their evolution the study will serve as a template for studying these other cratons.



Source: IE

PDF Reference URL: <https://www.drishtias.com/printpdf/emergence-of-earliest-continental-landmass>