

# Lassa fever

### Why in News?

Recently, three persons diagnosed with **Lassa fever** in the UK died. The cases have been linked to **travel** to west African countries.

### What is Lassa Fever?

#### About:

- The Lassa fever-causing virus is found in West Africa and was first discovered in 1969 in Lassa, Nigeria.
- The fever is spread by rats and is primarily found in countries in West Africa including Sierra Leone, Liberia, Guinea, and Nigeria where it is endemic.
  - Matomys rats has the potential to spread the deadly Lassa virus.
- The death rate associated with this disease is low, at around 1%. But the death rate
  is higher for certain individuals, such as pregnant women in their third trimester.
- According to the European Centre for Disease Prevention and Control, about 80% of the cases are asymptomatic and therefore remain undiagnosed.

#### Transmission:

- A person can become infected if they come in contact with household items of food that is contaminated with the urine or feces of an infected rat (zoonotic disease).
- It can also be spread, though rarely, if a person comes in contact with a sick person's
  infected bodily fluids or through mucous membranes such as the eyes, nose or the
  mouth.
  - Person-to-person transmission is more common in healthcare settings.

## Symptoms:

- Mild symptoms include slight fever, fatigue, weakness and headache.
- Serious symptoms include bleeding, difficulty breathing, vomiting, facial swelling, pain in the chest, back, and abdomen and shock.
- Death can occur from two weeks of the onset of symptoms, usually as a result of multiorgan failure.

#### Treatment:

• The **antiviral drug ribavirin** seems to be an effective treatment for Lassa fever if given early on in the course of clinical illness.

### Source: IE

