

Tropical Cyclones Need New Category

For Prelims: Tropical Cyclones Need New Category, <u>Tropical Cyclones</u>, Hurricanes, Saffir-Simpson (SS) Scale, <u>Global Warming</u>.

For Mains: Tropical Cyclones Need New Category, Important Geophysical phenomena such as earthquakes.

Source: TH

Why in News?

Recently, a study has been published in the journal **Proceedings of National Academy of Sciences**, where researchers have claimed that wind speed during a hurricane can cross 309 km/hour and therefore wind scale must add a Category 6.

What are the Key Highlights of the Study?

- Reconsideration of Saffir-Simpson (SS) Scale:
 - There are concerns about the adequacy of the Saffir-Simpson (SS) Hurricane Wind Scale, which has been used for over 50 years to communicate hurricane risk based solely on wind speed.
 - There are five categories on the SS hurricane wind scale category 1 to category 5 with category 5 wind speed exceeding 252 km/hour.
 - The combined effects of wind, storm surge, and rainfall in a category 5 impact would completely raze any structure.
 - The open-ended Category 5 may no longer be sufficient to communicate the increasing risk of hurricane damage in a warming climate.
- Introduction of Hypothetical Category 6:
 - Due to Global Warming, there is now a need to define a category 6 cyclone.
 - The warming can be observed not only at the sea surface, but also in the depths of the ocean, which increases the heat content of the ocean and thus favours the intensification of tropical cyclones.
 - To address the limitations of the existing scale the introduction of a hypothetical Category
 6 to the Saffir-Simpson Wind Scale is proposed with the wind speed above 309
 km/hour.
- Impact of Global Warming on Hurricane Intensification:
 - Increased <u>greenhouse gas emissions</u> have caused the Earth to warm by about 1.10 degrees Celsius since pre-industrial times and caused more intense tropical cyclones in the oceans.
 - For every degree of warming, the strongest cyclones are getting 12% stronger, making them 40% more destructive.

- As the oceans warm, cyclones also strengthen faster and spend more lifetime over the oceans.
 - In 2023, **tropical cyclone Freddy spent 37 days over the oceans,** making it the longest-lived cyclones ever recorded.
- Implications for Risk Messaging:
 - The findings underscore the **importance of revising risk messaging** to better inform the public about the increased risk of major hurricanes due to global warming.
 - SS Scale does not address issues related to inland flooding and storm surge, which are also critical components of hurricane risk.
 - Therefore, changes in messaging beyond wind-based scales are necessary to adequately communicate the full spectrum of hurricane hazards.

Note:

- Once a tropical cyclone reaches maximum sustained winds of 119 km/hour or higher, it is then
 classified as a hurricane, typhoon, or tropical cyclone, depending upon where the storm
 originates in the world.
 - In the North Atlantic, central North Pacific, and eastern North Pacific, the term hurricane is
- The Western Pacific basin is the most active region for tropical cyclones and accounts for about a third of the world's tropical cyclones.
- The North Indian basin accounts for only about 4% of the global total, although it is one of the most vulnerable regions in the world to the effects of such cyclones.



CYCLONE

Drishti IAS

Cyclones are rapid **inward** air circulation around a **low-pressure** area.




Cyclone v/s Anticyclone

Pressure System	Pressure Condition at the Center	Pattern of Wind Direction	
		Northern Hemisphere	Southern Hemisphere
Cyclone	ish'Low DishilAS	Anticlockwise	Clockwise
Anticyclone	High	Clockwise	Anticlockwise

Classification

- Tropical Cyclones; originate between the Tropics of Capricorn and Cancer
- Extra Tropical/ Temperate Cyclones; originate in the Polar Regions



Conditions for Formation

- Large sea surface with temperature >27° C.
- Presence of the Coriolis force
- · Small variations in the vertical wind speed
- · A pre-existing weak low- pressure area
- Upper divergence above the sea level system

Different Names for Tropical Cyclones


- Typhoons Southeast Asia and China
- Hurricanes North Atlantic and eastern Pacific
- Tornados West Africa and southern USA
- Willy-willies Northwest Australia
- Tropical Cyclones Southwest Pacific and Indian Ocean

Nomenclature

- Nodal Authority World Meteorological Organization (WMO)
- Indian Ocean Region Bangladesh, India, Maldives, Myanmar, Oman, Pakistan, Sri Lanka and Thailand contribute to naming cyclones that occur in this region.

Cyclones in India

- Bi-annual Cyclone Season March to May and October to December
- Recent Cyclones Tauktae, Vayu, Nisarga and Mekanu (in Arabian Sea) and Asani, Amphan, Fani, Nivar, Bulbul, Titli, Yaas and Sitrang (in Bay of Bengal)



Drishti Mains Question:

Q: Examine the limitations of the current Saffir-Simpson (SS) Scale and elucidate how the introduction of Category 6 can address these limitations.

UPSC Civil Services Examination Previous Year Question (PYQ)

Prelims

Q. Consider the following statements: (2020)

- 1. Jet streams occur in the Northern Hemisphere only.
- 2. Only some cyclones develop an eye.
- 3. The temperature inside the eye of a cyclone is nearly 10°C lesser than that of the surroundings.

Which of the statements given above is/are correct?

- (a) 1 only
- **(b)** 2 and 3 only
- (c) 2 only
- (d) 1 and 3 only

Ans: (c)

Q. In the South Atlantic and South-Eastern Pacific regions in tropical latitudes, cyclone does not originate. What is the reason? (2015)

- (a) Sea surface temperatures are low
- **(b)** Inter-Tropical Convergence Zone seldom occurs
- (c) Coriolis force is too weak
- (d) Absence of land in those regions

Ans: (b)

Mains:

Q. Tropical cyclones are largely confined to the South China Sea, Bay of Bengal and Gulf of Mexico. Why? **(2014)**

PDF Refernece URL: https://www.drishtiias.com/printpdf/tropical-cyclones-need-new-category