

Marmots

Why in News

Recently, reports of an outbreak of **bubonic plague** in Mongolia, China and far east Russia have emerged, caused mainly by **Tarbagan Marmot** (a **species of Marmot**).

■ It has been compared to the <u>Covid-19</u> <u>pandemic</u> which was apparently spread by the consumption of **bat** meat.

Key Points_//



General Description:

- Marmot (genus Marmota) belongs to the squirrel family (Sciuridae) within the order Rodentia.
- These have almost 15 species and the closest living relatives of marmots are ground squirrels and prairie dogs.
 - Prairie dogs are herbivorous burrowing rodents native to the grasslands of North America.
- Marmots are well suited for life in cold environments and have small fur-covered ears,
 short, stocky legs, and strong claws for digging.
- Marmots are diurnal (active during the day) and are almost entirely vegetarian.

Habitat:

- They are found primarily in the continents of **Europe, Asia and North America.**
- South Asia or the Indian Subcontinent is home to the Himalayan Marmot and the Long-tailed Marmot (both are Least Concerned in the <u>IUCN Red List</u>).
- Tarbagan or Mongolian Marmot (Endangered) is found in Mongolia, China and parts of Russia.

Importance:

- While digging burrows, marmots increase aeration in the soil which increases nutrient circulation and helps different plants to propagate.
- Marmots also play a very important role in Himalayan ecology. They are prey species for predators including the <u>snow leopard</u>, red fox, hawks, eagles, etc.

Threats:

- Marmots are hunted for their meat in China and Mongolia.
 - High altitude regions lack proteins. Pastoral nomads usually eat these when they do not have any other means of sustenance.
- Marmots are also hunted for their fur.

Plague

- Plague is caused by the bacteria Yersinia pestis usually found in small mammals and their fleas.
 - It is transmitted between animals and humans by the **bite of infected fleas, direct contact** with infected tissues and **inhalation of infected respiratory droplets.**
- It is one of the examples of <u>bacterial zoonoses</u>.
- There are **two main clinical forms** of plague infection:
 - Bubonic plague is the most common form and is characterized by painful swollen lymph nodes or 'buboes'.
 - Highly infectious bubonic plague killed about 50 million people across Africa, Asia and Europe in the 14th century.
 - Over 3,200 people were infected worldwide between 2000-15, resulting in 584 deaths.
 - The bacterial disease was named the Black Death after the dark swellings or buboes that victims suffered.
 - **Pneumonic plague** is a form of severe lung infection.
- Antibiotic treatment is effective against plague bacteria, so early diagnosis and early treatment can save lives. However, if left untreated, the fever can kill a victim in a very short time.

Source: DTE

PDF Reference URL: https://www.drishtiias.com/printpdf/marmots