

Vibrio Vulnificus Infection

Source: DTE

Why in News?


In recent years, India has been grappling with a growing concern related to **Vibrio vulnificus** infections, a deadly bacteria found in marine environments.

• Despite its potential threat, this pathogen remains largely underreported in India.

What is Vibrio Vulnificus?

- About:
 - Vibrio vulnificus is a bacterium that can cause severe infections in humans. It can result from eating undercooked seafood, especially oysters, which may contain the bacteria.
- Carriers:
 - It is typically contracted through two main routes: consuming infected raw shellfish and exposing wounds to contaminated waters.
 - It spreads through marine organisms like fish like eel, derbio, tilapia, trout, and shrimp.
 - The first case in marine organisms was documented in Japanese eel in 1975.

 The first case of V vulnificus in humans was recorded in 1976 in the US.
 - The pathogen arrived in Spain through imported eels in 1985.
 - In 2018, India documented an outbreak of V vulnificus in a tilapia farm in Kerala.
 - Originally from Africa and West Asia, <u>tilapia</u> is one of the most traded food fish globally.
- Symptoms:
 - Symptoms of V. vulnificus infection include diarrhea, vomiting, fever, and, in severe cases, flesh-eating diseases that can be fatal within days.
- Environmental Factors Favoring V. vulnificus in India:
 - This bacterium thrives in warm waters above 20°C. India's average sea surface temperature of 28°C provides a perfect habitat.
 - <u>Climate change</u>, with increased rainfall and reduced coastal salinity, further supports the growth of V. vulnificus.
- Consequences:
 - V. vulnificus infections have a **high mortality rate, ranging from 15% to 50%**, even with prompt diagnosis and treatment.
 - Vulnerable populations, such as those with chronic liver disease, cancer, chronic kidney disease, and diabetes, are at increased risk.
 - Infections can lead to **limb amputations** (surgical removal of part of the body, such as an arm or leg), making them a significant health concern.
- The Global Spread: //



Measures to Mitigate V. vulnificus Risk:

- Healthcare Awareness: Ensure that healthcare professionals in coastal areas are aware of V. vulnificus risks and test patients with relevant symptoms.
- Predictive Tools: Researchers are developing risk-warning tools using satellite-based sensors to monitor sea surface temperature and phytoplankton levels, which are associated with increased V. vulnificus infections.
- Learning from Seasonal Consumption in Japan: In Japan, bivalves like oysters and mussels are consumed only in winter, avoiding the summer when bacteria levels are high. This practice significantly reduces infection risk.

PDF Reference URL: https://www.drishtiias.com/printpdf/vibrio-vulnificus-infection