

Avian Botulism in Rajasthan

Why in News?

Recently, the <u>Centre for Avian Research Institute</u> reported the death of at least 600 migratory birds in Rajasthan.

• **High temperatures and reduced salinity in** <u>Sambhar Lake</u> likely created conditions that triggered <u>Avian Botulism</u>, causing the **mass deaths of** <u>migratory birds.</u>

Key Points

About Avian botulism:

- It is a <u>neuro-muscular illness</u> caused by **Botulinum (natural toxin**) that is produced by a bacteria Clostridium botulinum.
 - The bacteria is commonly found in the soil, rivers, and seawater. It affects both humans and animals.
 - It also needs anaerobic (absence of oxygen) conditions and does not grow in acidic conditions.
- It affects the nervous system of birds, leading to paralysis in their legs and wings.
 - Bacterial spores are widespread in wetland sediments and are commonly found in wetland habitats.
 - They are **present in invertebrates** like insects, mollusks, crustaceans, and even healthy vertebrates, including birds.
- The outbreaks of avian botulism tend to occur when average temperatures are above 21 degrees celsius, and during droughts.
- The deaths began on 26th October 2024, and continued for approximately two weeks.

Contributing Environmental Factors:

- Jaipur district, 70 km from Sambhar Lake, recorded above-average temperatures throughout October.
- Sambhar Lake experienced reduced oxygen levels due to the absence of rainfall.

Vulnerability of Migratory Birds

- Migratory birds arrive weakened from long journeys, making them more susceptible to diseases.
- Decaying bird carcasses attract maggots, which further contaminate the water and infect other birds or animals.

Management and Challenges

- Avian botulism cannot be treated, but immediate removal and disposal of affected birds are recommended to limit the spread.
- Sambhar Lake experienced a similar die-off in 2019, resulting in the deaths of nearly 18,000 birds.
- **Outbreaks are hard to predict** as they depend on specific environmental conditions aligning, such as a shift from high to low salinity coinciding with the arrival of migratory birds.


Global Perspective

- Spores of Clostridium botulinum can survive for years but produce toxins only under favorable environmental conditions.
- o Similar outbreaks have been observed in Australia and the United States during

periods of reduced salinity.

 Globally, around 57 diseases have been reported among wild birds, highlighting the broader ecological risks.

Sambhar Lake





tion:

• Situated about 80 km southwest of laipur, in east-central Rajasthan.

Features:

- It is the largest inland salt lake in India. It represents the depression of the Aravalli Range.
- The lake's salt supply was worked by the Mughal dynasty (1526-1857) and it
 was later owned jointly by the Jaipur and Jodhpur princely states.

Ramsar Site:

• It is a **wetland** of '**international importance'** under the <u>Ramsar Convention</u>, declared in 1990.

Rivers:

 It receives water from six rivers, namely Samaod, Khari, Mantha, Khandela, Medtha, and Roopangarh.

Vegetation:

- The vegetation present in the catchment area is mostly xerophytic type.
- Xerophyte is a plant adapted for growth under dry conditions.

The Central Avian Research Institute of India (CARI)

- It is a research institute located at Izzatnagar near Bareilly, Uttar Pradesh.
- It was established in 1979 under the administrative control of Indian Council of Agricultural Research (ICAR).
- It **studies poultry science**, including avian genetics, breeding, nutrition and feed technology, and avian physiology and reproduction, for the betterment of the Indian poultry industry.

