

Astronomers Uncover Hot Helium Stars

Source: TH

Why in News?

Astronomers have recently identified a group of **hot, helium-covered stars found in** <u>binary systems</u>, potentially deepening our understanding of stellar dynamics and evolution.

What are the Major Outcomes of the Discovery?

- Utilising a telescope capable of detecting <u>ultraviolet light</u>, astronomers observed around half a million stars in the <u>Large and Small Magellanic Clouds</u>.
 - The Large Magellanic Cloud and the Small Magellanic Cloud are two dwarf galaxies that are companions to the Milky Way.
- Some stars exhibited unusual speeds, hinting at the presence of companions (binary nature) affecting their motion.
 - Subsequent analysis of the optical spectra of 25 stars revealed their elemental composition, leading to the identification of different star classes.
- Stars were categorised into three classes:
 - Class 1 (helium-rich, hydrogen-depleted)
 - Class 2 and Class 3 (helium-rich, with hydrogen)


What is the Binary System of Stars?

- About: It refers to the pair of stars that are gravitationally bound to each other and orbit around a common centre of mass.
 - An estimated 85% or more of stars are actually part of binary or even multiple-star systems.
- Classification:
 - Visual Binaries: These are the easiest to identify and consist of two stars that can be directly resolved and separated using a telescope.
 - Spectroscopic binaries: These stars are too close together to be resolved visually even with powerful telescopes.
 - However, their presence can be detected by observing periodic shifts in their spectral lines.
 - **Eclipsing Binaries:** These binary systems are aligned in a way that one star periodically passes in front of the other from our perspective.
 - This event creates a **temporary dip in the brightness of the combined system**, allowing astronomers to confirm the presence of the unseen companion and study its properties.

- **Astrometric Binaries:** These binary systems are detected indirectly by measuring the **wobbling motion** of a single star.
 - This wobbling is caused by the gravitational pull of the unseen companion star.
- Confirmation of Binary Systems: When a star exhausts its fuel, gravity takes over, leading to a supernova explosion that strips its outer layers.
 - Some supernova lack **hydrogen**, suggesting pre-explosion stripping of the outer layer.
 - This can happen in binary systems, where **one star's gravity removes the outer hydrogen layer from its companion**, leaving behind a helium-rich star.
 - Astronomers have only found one such binary system so far.

//

Binary Star Orbit



How do Stars Maintain their Presence Over Billions of Years?

- Stars maintain their presence over billions of years through a delicate balance between two opposing forces: <u>nuclear fusion</u> and <u>gravity</u>.
- For example, despite **Newton's Law of Universal Gravitation** suggesting the Sun's gravitational collapse, nuclear fusion at its core acts as a vital stabilising force.
 - Nuclear Fusion involves the merging of nuclei of light elements like <u>hydrogen and helium</u>, releasing substantial <u>heat energy</u>.
 - This energy, in turn, creates internal pressure, counteracting the gravitational force, thus maintaining equilibrium.
- Therefore, Stars like the **Sun sustain this balance between outward fusion energy and inward gravitational pull**, ensuring their enduring presence over billions of years.

UPSC Civil Services Examination, Previous Year Question (PYQ)

Prelims:

- Q. The terms 'Event Horizon', 'Singularity', 'String Theory' and 'Standard Model' are sometimes seen in the news in the context of (2017)
- (a) Observation and understanding of the Universe

- (b) Study of the solar and the lunar eclipses
- (c) Placing satellites in the orbit of the Earth
- (d) Origin and evolution of living organisms on the Earth

Ans: (a)

Mains:

Q. How does the Juno Mission of NASA help to understand the origin and evolution of the Earth? (2017)

