



Reevaluating Obesity Metrics

[Source: HT](#)

The long-standing reliance on [Body Mass Index \(BMI\)](#) for diagnosing [obesity](#) is increasingly being questioned due to its **limitations**.

- BMI can **overestimate** obesity in **muscular individuals** like athletes and **underestimate** obesity in those with excessive fat but **low muscle mass**.
- [The Lancet](#) recommends using alternative metrics such as **waist circumference**, **waist-hip ratio**, and **waist-height ratio**, which consider **gender, age, and ethnicity** differences.
 - Obesity should be classified as **pre-clinical (no organ dysfunction)** and **clinical** (with **organ dysfunction** and activity impairment).
- **BMI** is a measure used to **assess** whether a person has a **healthy body weight** for a given **height**. It is calculated using a person's weight and height.
- **Obesity in India:** As per [The Lancet](#), **70%** of India's **urban population** is classified as **obese or overweight**.
 - **India ranks 3rd**, just behind the **US and China**, in the list of top 10 countries with the **highest number** of obese individuals.
 - Obesity is a health condition caused by **excessive body fat**, diagnosed when **BMI is 30 or higher**.

Read More: [Growing Obesity in India](#)

PDF Reference URL: <https://www.drishtias.com/printpdf/reevaluating-obesity-metrics>